

Microplastic abundance in beach sediments of the Kiel Fjord, Western Baltic Sea

Journal: Environmental Science and Pollution Research

Kevin Schröder ^a, Elke Kosse^b; Mark Lenz^b

- a. University of Kiel, Faculty of Agricultural and Nutritional Sciences, Olshausenstr. 40, 24098 Kiel, Germany
- b. GEOMAR Helmholtz Centre for Ocean Research Kiel, Marine Ecology, Düsternbrooker Weg 20, 24105 Kiel, Germany
- c. GEOMAR Helmholtz Centre for Ocean Research Kiel, Marine Biogeochemistry, Wischhofstr. 1 – 3, 24148 Kiel, Germany

Corresponding author: Mark Lenz, mlenz@geomar.de, Fon: ++49 431 600 4576, Fax: ++49 431 600 1671




Figure 1 Grain size analysis results for five sediment samples at the sewage treatment plant Bülk (BK1 – BK5). On the x-axis the grain size is displayed on a logarithmic scale. The y-axis displays the percentage share of the whole sediment sample




Figure 2 Grain size analysis results for five sediment samples at Falckenstein beach (FS1 – FS5). On the x-axis the grain size is displayed on a logarithmic scale. The y-axis displays the percentage share of the whole sediment sample




Figure 3 Grain size analysis results for five sediment samples at Tirpitzmole (TM1 –TM5). On the x-axis the grain size is displayed on a logarithmic scale. The y-axis displays the percentage share of the whole sediment sample